

KKU-IENC2014

"Engineering and Technological Responses to Global Challenges"



March 27 - 29, 2014
Pullman Khon Kaen Raja Orchid Hotel
Khon Kaen, Thailand

Abstract Collection



ANWAR
The 50th Anniversary of Khon Kaen University
and Faculty of Engineering

The 5th KKU International Engineering Conference 2014



Agricultural Machinery and
Postharvest Technology Center



Mercury Emission from Artisanal Buladu Gold Mine and Its Bioaccumulation in Rice Grains, Gorontalo Province, Indonesia

Anwar Mallongi^{1,a*}, Poranee Pataranawat^{2,b*}, Preeda Parkpian^{3,c},

¹ Postgraduate Education in Environmental Technology, Department of Sanitary Engineering, Faculty of Public Health, Mahidol University, Thailand and Faculty of Public Health, Hasanuddin University, Indonesia

² Department of Sanitary Engineering, Faculty of Public Health, Mahidol University, Thailand

³ School of Environment, Resources and Development, Asian Institute of Technology (AIT), Pathumthani, Thailand

⁴ Faculty of Public Health, Hasanuddin University ^aanwar_envi@yahoo.com,

^bpat_poranee@yahoo.com,

Keywords: Artisanal gold mine, THg mass balance, amalgamation, open burning, dry deposit, surface soil, rice grains

Abstract. Available forms of mercury (Hg) released from artisanal gold mine activities could be taken up increasingly by plants via root and leaf stomata. Total mercury (THg) concentrations in dry deposit, surface soil and rice grains were investigated as well as the potential risks in September 2011 from three rice fields of concern. The results revealed that the concentrations of THg in dry deposition, top soil (0 to 5 cm depth), sub soil (6 to 10 cm depth) and rice grains (*Oriza sativa* L.) both brown (once milled) and white (twice milled) grains were ranged from 166 to 322 $\mu\text{g m}^{-2} \text{ day}^{-1}$, 484 to 4244 $\mu\text{g kg}^{-1} \text{ dw}$, 122 to 1812 $\mu\text{g kg}^{-1} \text{ ww}$, and 113 to 1084 $\mu\text{g kg}^{-1} \text{ ww}$, respectively. Hazard quotient (HQ) values for dry deposition, top soil and sub soil were ranged from 3 to 7, 5 to 42 and 5 to 36, respectively. Target hazard quotient (THQ) for brown and white rice grains consumptions were found in the range of 0.1 to 1.6 and 0.1 to 1.0, respectively. THQ values through brown rice consumption exceeded the guideline (>1) presented that the brown rice in these areas should not be safe for consumption and are at risks for the whole lifespan. However, the THQ values of both brown and white rice grains in some areas were still low and should be safe for the whole life span consumption.